



VENTILATOR ASSOCIATED PNEUMONIA

Practice Alert Statements:

- ☑ All patients receiving mechanical ventilation, as well as those at high risk for aspiration (e.g., decreased level of consciousness; enteral tube in place), should have the head of the bed (HOB) elevated at an angle of 30 to 45° unless medically contraindicated.¹⁻⁷ (Level VI)
- ☑ Use an endotracheal tube (ET) with a dorsal lumen above the endotracheal cuff to allow drainage by continuous suctioning of tracheal secretions that accumulate in the subglottic area.^{1,2,8-13} (Level VI)
- ☑ Do not routinely change, on the basis of duration of use, the patient's ventilator circuit.^{1,14-17} (Level VI)

Supporting Evidence:

- Critically ill patients who are intubated for > 24 hours are at 6 to 21 times the risk of developing ventilator-associated pneumonia (VAP)^{1,2,18-20} and those intubated for < 24 hours are at 3 times the risk of VAP.²⁰ Other risk factors for VAP include decreased level of consciousness, supine positioning with HOB flat, use of H₂ antagonists and antacids, gastric distention, presence of gastric or small intestine tubes, enteral feedings, and a trauma or COPD diagnosis.^{1,18-22} VAP is reported to occur at rates of 10 to 35 cases / 1000 ventilator days, depending on the clinical situation.^{1,19}
- Morbidity and mortality associated with the development of VAP is high, with mortality rates ranging from 20 to 41%.^{20,23-25} Development of VAP increases ventilator days, critical care and hospital lengths of stay (LOS) by 4, 4 and 9 days, respectively,^{18,23,26} and results in > \$11,000 additional costs / VAP case.^{18,25,27}
- Micro or macro aspiration of oropharyngeal and/or gastric fluids are presumed to be an essential step in the development of VAP.^{1,2,12,28} Pulmonary aspiration is increased by supine positioning and pooling of secretions above the ET tube cuff.^{1,3,19}
- Compared to supine positioning, studies have shown that simple positioning with HOB elevation to 30° or higher significantly reduces gastric reflux and VAP,³⁻⁷ yet national surveys and reports in the literature describe poor compliance rates with HOB elevation in critical care units.^{20,29-34}
- Studies on the use of special ET tubes which remove secretions pooled above the cuff with continuous suction decrease VAP by 45 to 50 %.⁸⁻¹¹
- Studies on the frequency of ventilator circuit changes have found no increase in VAP with prolonged use.¹⁴⁻¹⁷
- National regulatory and expert consensus groups include the AACN VAP Practice Alert interventions as critical to decreasing VAP rates.^{1,2,35-37}

AACN Grading Level of Evidence

- Level I. Manufacturers recommendation only.
- Level II. Theory based-no research data to support recommendations; Recommendations from expert consensus group may exist.
- Level III. Laboratory or bench data only-no clinical data to support recommendations
- Level IV. Limited clinical studies to support recommendations.
- Level V. Clinical studies in more that one or two different populations or situations to support recommendations.
- Level VI. Clinical studies in a variety of patient populations and situations to support recommendations.

Actions for Nursing Practice:

- Always keep mechanically ventilated patients HOB elevated to 30° or higher, unless medically contraindicated; use an ET tube with continuous suction above the cuff in patients expected to be intubated > 48 hours; do not routinely change ventilator circuits.
- Assure that your critical care unit has written practice documents such as a policy, procedure or standards of care that includes these practice alerts.³⁸

- Determine your unit's rate of compliance with the HOB elevation directive and use an ET tube with continuous suction above the cuff ([click here for PI audit tool](#))
- If compliance with HOB elevation is < 90%, develop a plan to improve compliance^{29,38-45}
 - ⇒ Consider forming a multidisciplinary task force (nurses, physicians, respiratory therapists, clinical pharmacists) to address VAP practice changes.
 - ⇒ Educate staff about the significance of hospital-acquired pneumonias in critically ill patients and how the interventions listed in the Practice Alert can reduce VAP ([click for education slide program](#)).
 - ⇒ Incorporate content into orientation programs, initial and annual competency verifications.
 - ⇒ Develop a variety of communication strategies to alert and remind staff of the importance of these VAP interventions and to disseminate results of audits.
 - ⇒ Develop documentation standards for HOB elevation that include a rationale for when HOB elevation is not done.
 - ⇒ Incorporate HOB elevation to at least 30° in mechanically ventilated patients or patients at high risk for aspiration in any unit standing orders. Also include HOB elevation monitoring in your critical care scorecard / QI plan / PI activities to assure that compliance levels are maintained.

Expected Outcomes:

- Decrease in VAP rates for the unit
- Increase in number of patients with HOB elevation to at least 30°
- Cost savings due to decreased rates of VAP and less frequent ventilator circuit changes

Resources:

- Education Materials:
 - ⇒ Power Point slide program for VAP education sessions (www.aacn.org)
 - ⇒ Online continuing education program on prevention strategies for VAP (<http://www.nellcor.com/educ/onlineed.aspx>)
- Audit Tools:
 - ⇒ Measurement of compliance with HOB elevation in mechanically ventilated patients (www.aacn.org)
- Others:
 - ⇒ Methods for estimating HOB elevation (www.aacn.org)
 - ⇒ For additional information/assistance, contact a clinical practice specialist with the AACN Practice Resource Network (PRN) via email at practice@aacn.org or via phone at 1-800-394-5995 x217.

References:

1. Tablan OC, Anderson LJ, Besser R, Bridges C, Hajjeh R; CDC; Healthcare Infection Control Practices Advisory Committee. Guidelines for preventing health-care associated pneumonia, 2003: Recommendations of CDC and the Healthcare Infection Control Practices Advisory Committee. *Morbidity Mortality Weekly Review* (MMWR). 2004;53(RR-3):1-36. <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5303a1.htm> (accessed 2/12/07).
2. American Thoracic Society; Infectious Diseases Society of America. Guidelines for the management of adults with hospital-acquired, ventilator-associated, and healthcare-associated pneumonia. *American Journal Respiratory Critical Care Medicine*. 2005;171:388-416.
3. Torres A, Serra-Batiles J, Ros E, et al. Pulmonary aspiration of gastric contents in patients receiving mechanical ventilation: the effect of body position. *Annals Internal Medicine* 1992;116:540-542.
4. Ibanez J, Penafiel A, Raurich J, et al. Gastroesophageal reflux in intubated patients receiving enteral nutrition: effect of supine and semi recumbent positions. *Journal Parenteral Enteral Nutrition*. 1992;16:419-422.
5. Orozco-Levi M, Torres A, Ferrer M, et al. Semi-recumbent position protects from pulmonary aspiration but not completely from gastroesophageal reflux in mechanically ventilated patients. *American Journal Respiratory Critical Care Medicine*. 1995;152:1387-1390.
6. Drakulovic M, Torres A, Bauer T, et al. Supine body position as a risk factor for nosocomial pneumonia in mechanically ventilated patients: A randomized trial. *Lancet*. 1999;354:1851-1854.
7. Davis K Jr, Johannigman J, Campbell R, et al. The acute effects of body position strategies and respiratory therapy in paralyzed patients with acute lung injury. *Critical Care*. 2001;5:81-87.
8. Valles J, Artigas A, Rello J, et al. Continuous aspiration of subglottic secretions in preventing ventilator-associated pneumonia. *Intensive Care Medicine*. 1995;122:179-186.
9. Mahul P, Auboyer C, Jospe R, et al. Prevention of nosocomial pneumonia in Intubated patients: respective role of mechanical subglottic secretion drainage and stress ulcer prophylaxis. *Intensive Care Medicine*. 1992;18:20-25.
10. Kollef M, Skubas N, Sundt T. A randomized clinical trial of continuous aspiration of subglottic secretions in cardiac surgery patients. *Chest*. 1999;116:1339-1346.

11. Dezfulian C, Shojania K, Collard HR, Kim HM, Matthay MA, Saint S. Subglottic secretion drainage for preventing ventilator-associated pneumonia: a meta-analysis. *American Journal Medicine*. 2005;118:11-8.
12. Cook D, KeJonge B, Brochard L, Brun-Buisson C. Influence of airway management on ventilator-associated pneumonia: evidence from randomized trials. *Journal American Medical Association*. 1998;279:761-787.
13. Smulders K, van der Hoeven H, Weers-Pothoff I, Vandernbroucke-Grauls C. A randomized clinical trial of intermittent subglottic secretion drainage in patients receiving mechanical ventilation. *Chest*. 2002;121:858-862.
14. Dreyfuss D, Djedaini K, Weber P, et al. Prospective study of nosocomial pneumonia and of patient circuit colonization during mechanical ventilation with circuit changes every 48 hours versus no change. *American Journal Respiratory Critical Care Medicine*. 1991;143:738-743.
15. Kotilainen H, Keroack M. Cost analysis and clinical impact of weekly ventilator circuit changes in patients in intensive care unit. *American Journal Infection Control*. 1997;25:117-120.
16. Kollef M, Shapiro S, Fraser V, et al. Mechanical ventilation with or without 7-day circuit changes: a randomized controlled trial. *Annals Internal Medicine*. 1995;123:168-174.
17. Long M, Wickstrom G, Grivmes A, Benton C, Belcher B, Staff A. Prospective, randomized study of ventilator-associated pneumonia in patients with one versus three ventilator circuit changes per week. *Infection Control Hospital Epidemiology*. 1996;17:14-19.
18. Rello J, Ollendorf D, Oster G, et al. Epidemiology and outcomes of ventilator-associated pneumonia in a large US database. *Chest*. 2002;122:2115-2121.
19. Craven D. Epidemiology of ventilator-associated pneumonia. *Chest* 2000;117:186S-187S.
20. Kollef M. Ventilator-associated pneumonia: A multivariate analysis. *Journal American Medical Association*. 1993;270:1965-1970.
21. Flanders SA, Collard HR, Saint S. Nosocomial pneumonia: state of the science. *American Journal Infection Control*. 2006;34:84-93..
22. Elatrous S, Boujdaria R, Merghili S. Incidence and risk factors of ventilator-associated pneumonia: a one-year prospective study. *Clinical Intensive Care*. 1996;20:193-198.
23. Bercault N, Boulain T. Mortality rate attributable to ventilator-associated nosocomial pneumonia in an adult intensive care unit: A prospective case-control study. *Critical Care Medicine*. 2001;29:2303-2309.
24. Heyland D, Cook D, Griffith L, et al. The attributable morbidity and mortality of ventilator-associated pneumonia in the critically ill patient. *American Journal Respiratory Critical Care Medicine*. 1999;159:1249-1256.
25. Safdar N, Dezfulian C, Collard HR, Saint S. Clinical and economic consequences of ventilator-associated pneumonia: a systematic review. *Critical Care Medicine*. 2005;33:2184-93.
26. Chastre J, Fagon J. Ventilator-associated pneumonia. *American Journal of Respiratory Critical Care Medicine*. 2002;165:867-903.
27. Warren K, Shukla S, Olsen M, et al. Outcomes and attributable cost of ventilator-associated pneumonia among intensive care unit patients in a suburban medical center. *Critical Care Medicine*. 2003;31:1312-1317.
28. Bonten M, Kollef M, Hall J. Risk factors for ventilator-associated pneumonia: from epidemiology to patient management. *Healthcare Epidemiology*. 2004;38:1141-1149.
29. Zack J, Garrison T, Trouillion E, et al. Effect of an educational program aimed at reducing the occurrence of ventilator-associated pneumonia. *Critical Care Medicine* .2002;30:2407-2412.
30. Berenholtz S, Pronovost P. Barriers to translating evidence into practice. *Current Opinions Critical Care*. 2003;9:321-325.
31. Grap M, Cantly M, Munro C, et al. Use of backrest elevation in critical care: pilot study. *American Journal Critical Care*. 1999;8:475-480.
32. van Nieuwenhoven CA, Vandembroucke-Grauls C, van Tiel FH, et al. Feasibility and effects of the semirecumbent position to prevent ventilator-associated pneumonia: a randomized study. *Critical Care Medicine*. 2006;34:396-402.
33. Grap M, Munro C, Bryant S, Ashtiani B. Predictors of backrest elevation in critical care. *Intensive Critical Care Nursing*. 2003;19:68-74.
34. Grap MJ, Munro CL. Preventing ventilator-associated pneumonia: evidence-based care. *Crit Care Nurs Clin North Am*. 2004;16:349-358.
35. Joint Commission on Accreditation of Healthcare Organizations (JCAHO). *Specification Manual for National Hospital Quality Measures – ICU*. <http://www.jointcommission.org/PerformanceMeasurement/MeasureReserveLibrary/Spec+Manual+-+ICU.htm> (accessed February 12, 2007).
36. Burns S (editor). *AACN Protocols for Practice: Care of Mechanically Ventilated Patients, 2nd edition*. Jones and Bartlett Publishers:Sudbury, MA, 2007.
37. Shojania K, McDonald K, Wachter R, Owens D. *Closing the Quality Gap: A Critical Analysis of Quality Improvement Strategies. Volume 6 - Prevention of Healthcare-associated Infections*. Agency for Healthcare Research and Quality (AHRQ). 2007, page 71 to 105 (AHRQ publication No. 04(07)-0051-6). www.ahrq.gov/clinic/ptsafety/chap17a.htm. (accessed 2/12/2007).

38. Helman D, Sherner J, Fitzpatrick T, Callender M, Shorr A. Effect of standardized orders and provider education on head-of-bed positioning in mechanically ventilated patients. *Critical Care Medicine*. 2003;31:2285-2290.
39. Craven DE. Preventing ventilator-associated pneumonia in adults: sowing seeds of change. *Chest*. 2006;130(1):251-60.
40. Babcock H, Zack J, Carrison T et al. An educational intervention to reduce ventilator-associated pneumonia in an integrated health system: a comparison of effects. *Chest*. 2004;125:2224-2231.
41. Grap MJ, Munro CL. Quality improvement in backrest elevation: improving outcomes in critical care. *AACN Clinical Issues*. 2005;16(2):133-139.
42. Fox MY. Toward a zero VAP rate: personal and team approaches in the ICU. *Critical Care Nursing Quarterly*. 2006;29(2):108-14.
43. Murray T, Goodyear-Bruch C. Ventilator associated pneumonia improvement program. *AACN Advanced Critical Care*. 2007;18(2): in press.
44. Kollef M. Prevention of hospital-associated pneumonia and ventilator-associated pneumonia. *Critical Care Medicine*. 2004;32:1396-1405.
45. Isakow W, Kollef MH. Preventing ventilator-associated pneumonia: an evidence-based approach of modifiable risk factors. *Seminars Respiratory Critical Care Medicine*. 2006;27:5-17.